



**PERINATAL CARE
IN THE CONTEXT
OF COVID-19**

The KNH experience

BY JULIE MATIBA



Suspect cases

A. A patient with acute respiratory illness (fever or cough or shortness of breath) AND

- A history of travel to a foreign country during the 14 days prior to symptom onset, or
- Having been in contact with a confirmed or probable case of COVID-19 in the 14 days prior to symptom onset, or

B. A patient with severe acute respiratory illness (fever or cough or shortness of breath; AND requiring hospitalization)

And in the absence of alternative diagnosis that fully explains the clinical presentation

Probable case

- A suspect case for whom testing could not be performed for any reason OR
- A suspect case for whom testing for COVID-19 is inconclusive or for whom testing was positive on a pan-coronavirus assay.

Confirmed case

- A person with laboratory confirmation of COVID-19 infection, irrespective of clinical signs and symptoms

PPEs for Different Clinical Settings

	Outside OR				Emergent Tracheal intubation	Inside OR			
	Public Area	Outpatient Clinic, Ward	ER	Fever Clinic	Exception of Fever Clinic	Office, Cafeteria, locker room, etc.	Non-febrile patients	Febrile patients	Suspected or confirmed COVID-19 cases
Head	Disposable Surgical Cap								
Mask	Surgical mask	Surgical mask	Surgical mask	N95 mask	N95 mask	Surgical mask	Surgical mask	N95 mask	N95 mask
Glove	-	-	+	+	+	-	-	+	+
Shoe covers	-	-	+	+	+	-	-	+	+
Protective suit	-	-	-	+	-	-	-	-	+
Isolation gown	-	-	+	+	+	-	-	+	+
Goggles or protective shield	-	-	±	+	+	-	+	+	+




Figure 2. Enhanced traffic control bundling.